

UCC 504.5:635.8:664.8.022.7
DOI: 10.37128/2707-5826-2020-2-19

ANALYSIS OF THE EFFICIENCY OF USING THE CULINARY PROCESSING OF MUSHROOMS IN ORDER TO REDUCE THE CONCENTRATION OF HEAVY METALS IN THEM

O.I. ВРАДІЙ, асистент,
Вінницький національний
аграрний університет

The samples of mushrooms were collected in the forestry of the village Tyvriv and the town of Kalynivka, Vinnytsia and Kalynivka districts, in the conditions of Right-Bank Forest-Steppe of Ukraine. The studies of heavy metals concentration were performed in the scientific agrochemical laboratory of the Department of Ecology and Environmental Protection, the Faculty of Agronomy and Forestry at Vinnytsia National Agrarian University. The concentrations of Cu, Zn, Pb and Cd of the studied mushrooms were determined by the method of atomic absorption spectrometry after the dry mineralization. The research of the effect of culinary processing on the concentration of heavy metals in mushrooms were conducted according to the following scheme: 1. The control: mushrooms, cleaned and washed with tap water; 2. Option 1: mushrooms soaked in tap water for 3 hours and boiled in it; 3. Option 2: mushrooms soaked in the water without mineral residue for 3 hours and boiled in it; 4. Option 3: mushrooms soaked in distilled water for 3 hours and boiled in it.

Summarizing the obtained results, it should be noted that the most effective way to reduce the concentration of lead and cadmium in mushrooms was the option of soaking and boiling mushrooms in the water without mineral residue. The most effective way to reduce the concentration of zinc and copper was Option 3, where distilled water was used. The influence of soaking and boiling mushrooms in different types of water is investigated. When soaking and boiling mushrooms in tap water for 3 hours, the concentration of zinc in them was found to decrease from 1.01 to 1.04 times, the concentration of copper – from 1.0 to 1.2 times, the concentration of lead – from 1.09 to 1.16 times and the concentration of cadmium – from 1.08 to 1.2 times. When soaking and boiling mushrooms in the water without mineral residue, the concentration of zinc decreased from 1.01 to 1.04 times, the concentration of copper – from 1.16 to 1.25 times, the concentration of lead – from 1.8 to 2.0 times and the concentration of cadmium – from 1.75 to 2.0 times. When soaking mushrooms and boiling them in distilled water, the concentration of zinc decreased from 1.13 to 1.18 times, the concentration of copper – from 1.2 to 1.29 times, the concentration of lead – from 1.22 to 1.28 times and the concentration of cadmium – from 1.08 to 1.2 times.

Key words: mushrooms, concentration, zinc, copper, lead, cadmium, distilled water, water without mineral residue, tap water, boiling of mushrooms.

Tabl. 4. Fig. 4. Lit. 9.

Setting the problem. Mushrooms have always been considered a valuable food product, and mushroom dishes are a favorite food of many peoples. This is not accidental, since they contain many organic and mineral compounds, similar in chemical composition to vegetables and products of animal origin [1, 2]. It is known that fresh mushrooms contain: 84-94% of water, 2-6% of nitrogenous substances (80% of which are proteins), 1-3% of carbohydrates, 0.2-6.0% of fats, 0.6-1.0% of minerals, as well as vitamins A, B1, B2, C, D, PP [3-4].

The high content of chitin forms complexes with proteins, pigments and calcium salts. It differs significantly from vegetable fiber – cellulose and prevents the complete absorption of mushroom dishes during digestion. In the dry matter of mushrooms there are many proteins [5], the content of which often exceeds 40%, that is, it is higher than in the seeds of peas and beans. The digestibility of fungal proteins is 54-85%, that is, approximately the same as that of vegetable ones. The carbohydrate content in the dry matter of mushrooms can reach 70% and they are absorbed by 93-99%. Such carbohydrates as Glycogen (animal starch), Trehalose, Glucose, Mycosis and Mannitol are dominated in mushrooms. These substances cause a pleasant sweet taste of mushrooms. Many species are characterized by the high content of carbohydrates in the dry matter. They are still little used, but they are widespread and suitable for sautéing, in particular *Russula Pers.* (*Lat.*) and *Lactarius resimus* (*Lat.*). [7] Fatty substances of mushrooms (phosphatides, cholesterol and ergosterol) are absorbed by 92-97%, and their content in the fruiting bodies is greater than in potatoes or other vegetables. In the fruiting bodies there are also extractive substances, carotene, pantothenic acid, enzymes and antibiotic substances, in addition to various mineral substances, in the content of which mushrooms are close to fruit [8-9].

The forestry products of vegetable origin, namely their harvesting is of great importance for the enterprises of the forest industry, providing people with food of high value. In particular, mushrooms and berries sometimes supplement people's diets, but there are also occasions when they serve staple foods, even on a par with bread, meat and vegetables.

Wild berries and edible mushrooms are mostly used for consumption without special processing. That is why the sanitary and hygienic quality of non-timber raw materials is a significant factor affecting its safety [5]. However, the content of heavy metals in the phytomass of representatives of Ukraine's wild flora, suitable for using as non-timber raw materials, has not been sufficiently studied. The bulk of the research is devoted to the characteristics of migration and accumulation of heavy metals in crops, while wild species remain outside the focus of scientists. Because of the intensification of technogenic and anthropogenic pressure on ecosystems, the number of areas suitable for the collection of environmentally friendly non-timber raw materials is sharply reduced.

The consumption of non-timber forest resources containing heavy metals can lead to adverse effects occurring in the human body. Therefore, the topic of finding ways to reduce the content of these substances in forestry products is very relevant.

Analysis of recent researches and publications. The studies on the bioaccumulation of heavy metals in various types of phytocenoses are of considerable interest, since plants and mushrooms, as an autotrophic block of ecosystems, play an important role in the redistribution of chemical elements between individual components of the biosphere. The forest landscapes, as dominant geochemical regulators of cyclic mass flows of heavy metals at the global level, are of particular importance in this process [10-11].

The excessive concentrations of heavy metals in the environment, far exceeding the natural ones, can greatly affect the functioning of plants and mushrooms, reduce the biotic productivity of forests, impair their recreational potential and alter the quality of wood, which may affect its environmental safety. In the complex of researches on the biotic productivity, the study of the role of timber and non-timber forest resources in the biogeocenoses in relation to the migration and deposition of elements of the heavy metals group, especially in the area of technogenic loading, is an urgent and timely task. Its solution will allow finding the mechanisms of selective ability of forest species to absorb and accumulate heavy metals in tissues, and to remove them from circulation for a long time. The excessive concentrations of heavy metals can influence the course of physiological processes to some extent that is reflected in the change of the integral indicator of forest plants – phytomass. According to the literature, timber and non-timber forest resources respond to the environmental pollution by reducing their biotic productivity [12]. Therefore, the need to monitor the pollution of non-timber forest resources by heavy metals and to study the ways to improve their quality remains relevant.

Research methodology and conditions. The samples of mushrooms were collected in the forestry of the village Tyvriv and the town of Kalynivka, Vinnytsia and Kalynivka districts, in the conditions of Right-Bank Forest-Steppe of Ukraine. The studies of heavy metals concentration were performed in the scientific agrochemical laboratory of the Department of Ecology and Environmental Protection, the Faculty of Agronomy and Forestry at Vinnytsia National Agrarian University. The concentrations of Cu, Zn, Pb and Cd of the studied mushrooms were determined by the method of atomic absorption spectrometry after the dry mineralization [13, 14]. The research of the effect of culinary processing on the concentration of heavy metals in mushrooms were conducted according to the following scheme: 1. The control: mushrooms, cleaned and washed with tap water; 2. Option 1: mushrooms soaked in tap water for 3 hours and boiled in it; 3. Option 2: mushrooms soaked in the water without mineral residue for 3 hours and boiled in it; 4. Option 3: mushrooms soaked in distilled water for 3 hours and boiled in it.

The purpose of the article is to study the efficiency of using the methods of culinary processing of mushrooms in order to reduce the content of heavy metals in them.

Presenting the main material. The obtained results of the research showed a certain influence of soaking and boiling mushrooms in different types of water on the concentration of heavy metals in them (Table 1-4).

Table 1

Zinc concentration in mushrooms, mg/kg

Kind of mushrooms	MPC	The control	Option 1	Option 2	Option 3
<i>Cantharellus cibarius</i> (Lat.)	20	6.41	6.26	6.24	5.64
<i>Gyroporus cyanescens</i> (Lat.)	20	7.09	6.97	6.95	6.1
<i>Laetiporus sulphureus</i> (Lat.)	20	5.04	4.91	4.81	4.26
<i>Butyriboletus regius</i> (Lat.)	20	10.99	10.65	10.62	9.4
<i>Leccinum</i> (Lat.)	20	7.86	7.72	7.62	6.84
<i>Russula Pers.</i> (Lat.)	20	11.18	10.75	10.68	9.49
<i>Boletus edulis</i> (Lat.)	20	11.41	11.08	11.16	9.74
<i>Amanita rubescens</i> (Lat.)	20	6.59	6.35	6.32	5.78
<i>Leccinum scabrum</i> (Lat.)	20	4.16	4.09	4.01	3.5
<i>Leccinum aurantiacum</i> (Lat.)	20	10.32	9.99	9.83	8.86
<i>Armillaria mellea</i> (Lat.)	20	0.074	0.073	0.073	0.064

The source is based on our own research findings

In Option 1 (Table 1) the concentration of zinc in mushrooms decreased from 1.01 to 1.04 times, in particular, in *Cantharellus cibarius* (Lat.) – by 1.02 times, in *Gyroporus cyanescens* (Lat.) – by 1.01 times, in *Laetiporus sulphureus* (Lat.) – by 1.02 times, in *Butyriboletus regius* (Lat.) – by 1.03 times, in *Leccinum* (Lat.) – by 1.01 times, in *Russula Pers.* (Lat.) – by 1.04 times, in *Boletus edulis* (Lat.) – by 1.02 times, in *Amanita rubescens* (Lat.) – by 1.03 times, in *Leccinum scabrum* (Lat.) – by 1.01 times, in *Leccinum aurantiacum* (Lat.) – by 1.03 times and in *Armillaria mellea* (Lat.) – by 1.01 times.

In Option 2 the concentration of zinc in the mushrooms decreased from 1.01 to 1.04 times, in particular, in *Cantharellus cibarius* (Lat.) and *Gyroporus cyanescens* (Lat.) – by 1.02 times, in *Laetiporus sulphureus* (Lat.) – by 1.04 times, in *Butyriboletus regius* (Lat.) and *Leccinum* (Lat.) – by 1.03 times, in *Russula Pers.* (Lat.) – by 1.04 times, in *Boletus edulis* (Lat.) – by 1.02 times, in *Amanita rubescens* (Lat.) – by 1.04 times, in *Leccinum scabrum* (Lat.) – by 1.03 times, in *Leccinum aurantiacum* (Lat.) – by 1.04 times and in *Armillaria mellea* (Lat.) – by 1.01 times.

In Option 3 the concentration of zinc in mushrooms decreased from 1.13 to 1.18 times, in particular, in *Cantharellus cibarius* (Lat.) – by 1.13 times, in *Gyroporus cyanescens* (Lat.) – by 1.16 times, in *Laetiporus sulphureus* (Lat.) – by 1.18 times, in *Butyriboletus regius* (Lat.) – by 1.16 times, in *Leccinum* (Lat.) – by 1.14 times, in *Russula Pers.* (Lat.) and *Boletus edulis* (Lat.) – by 1.17 times, in *Amanita rubescens* (Lat.) – by 1.14 times, in *Leccinum scabrum* (Lat.) – by 1.18 times, in *Leccinum aurantiacum* (Lat.) – by 1.16 times and in *Armillaria mellea* (Lat.) – by 1.15 times (Fig. 1).




Fig. 1 Concentration of zinc in mushrooms

The source is based on our own research findings

In Option 1 (Table 2) the concentration of copper in mushrooms decreased from 1.0 to 1.2 times, in particular, in *Cantharellus cibarius* (Lat.) – by 1.03 times,

Table 2

Copper concentration in mushrooms, mg/kg

Kind of mushrooms	MPC	The control	Option 1	Option 2	Option 3
<i>Cantharellus cibarius</i> (Lat.)	10	0.32	0.31	0.26	0.25
<i>Gyroporus cyanescens</i> (Lat.)	10	0.63	0.62	0.51	0.49
<i>Laetiporus sulphureus</i> (Lat.)	10	0.06	0.05	0.05	0.05
<i>Butyriboletus regius</i> (Lat.)	10	0.18	0.17	0.15	0.14
<i>Leccinum</i> (Lat.)	10	0.25	0.24	0.2	0.2
<i>Russula Pers.</i> (Lat.)	10	0.64	0.62	0.51	0.51
<i>Boletus edulis</i> (Lat.)	10	0.26	0.25	0.21	0.21
<i>Amanita rubescens</i> (Lat.)	10	0.16	0.15	0.13	0.13
<i>Leccinum scabrum</i> (Lat.)	10	0.70	0.68	0.57	0.55
<i>Leccinum aurantiacum</i> (Lat.)	10	0.14	0.13	0.12	0.11
<i>Armillaria mellea</i> (Lat.)	10	2.80	2.72	2.25	2.17

The source is based on our own research findings

in *Gyroporus cyanescens* (Lat.) – by 1.01 times, in *Laetiporus sulphureus* (Lat.) – by 1.2 times, in *Butyriboletus regius* (Lat.) – by 1.05 times, in *Leccinum* (Lat.) – by 1.04 times, in *Russula Pers.* (Lat.) – by 1.0 times, in *Boletus edulis* (Lat.) – by 1.04 times, in *Amanita rubescens* (Lat.) – by 1.06 times, in *Leccinum scabrum* (Lat.) – by 1.02

times, in *Leccinum aurantiacum* (Lat.) – by 1.07 times and in *Armillaria mellea* (Lat.) – by 1.02 times.

In Option 2 the concentration of copper in mushrooms decreased from 1.16 to 1.25 times, in particular, in *Cantharellus cibarius* (Lat.) and *Gyroporus cyanescens* (Lat.) – by 1.23 times, in *Laetiporus sulphureus* (Lat.) and *Butyriboletus regius* (Lat.) – by 1.2 times, in *Leccinum* (Lat.) and *Russula Pers.* (Lat.) – by 1.25 times, in *Boletus edulis* (Lat.) and *Amanita rubescens* (Lat.) – by 1.23 times, in *Leccinum scabrum* (Lat.) – by 1.22 times, in *Leccinum aurantiacum* (Lat.) – by 1.16 times and in *Armillaria mellea* (Lat.) – by 1.24 times.

In Option 3 the concentration of copper in mushrooms decreased from 1.2 to 1.29 times, in particular, in *Cantharellus cibarius* (Lat.) and *Gyroporus cyanescens* (Lat.) – by 1.28 times, in *Laetiporus sulphureus* (Lat.) – by 1.2 times, in *Butyriboletus regius* (Lat.) – by 1.28 times, in *Leccinum* (Lat.) and *Russula Pers.* (Lat.) – by 1.25 times, in *Boletus edulis* (Lat.) and *Amanita rubescens* (Lat.) – by 1.23 times, in *Leccinum scabrum* (Lat.) and *Leccinum aurantiacum* (Lat.) – by 1.27 times, and in *Armillaria mellea* (Lat.) – by 1.29 times (Fig. 2).




Fig. 2. Concentration of copper in mushrooms

The source is based on our own research findings

Table 3

Lead concentration in mushrooms, mg/kg

Kind of mushrooms	MPC	The control	Option 1	Option 2	Option 3
<i>Cantharellus cibarius</i> (Lat.)	0.5	0.21	0.19	0.11	0.17
<i>Gyroporus cyanescens</i> (Lat.)	0.5	0.22	0.20	0.11	0.17
<i>Laetiporus sulphureus</i> (Lat.)	0.5	0.27	0.24	0.15	0.22
<i>Butyriboletus regius</i> (Lat.)	0.5	0.24	0.21	0.13.	0.19
<i>Leccinum</i> (Lat.)	0.5	0.28	0.25	0.15	0.22
<i>Russula Pers.</i> (Lat.)	0.5	0.21	0.18	0.11	0.17
<i>Boletus edulis</i> (Lat.)	0.5	0.23	0.21	0.12	0.18
<i>Amanita rubescens</i> (Lat.)	0.5	0.27	0.24	0.15	0.21
<i>Leccinum scabrum</i> (Lat.)	0.5	0.26	0.23	0.14	0.21
<i>Leccinum aurantiacum</i> (Lat.)	0.5	0.22	0.20	0.12	0.18
<i>Armillaria mellea</i> (Lat.)	0.5	0.29	0.26	0.16	0.23

The source is based on our own research findings

In Option 1 (Table 3) the concentration of lead in mushrooms decreased from 1.09 to 1.16 times, in particular, in *Cantharellus cibarius* (Lat.) – by 1.1 times, in *Gyroporus cyanescens* (Lat.) – by 1.1 times, in *Laetiporus sulphureus* (Lat.) – by 1.12 times, in *Butyriboletus regius* (Lat.) – by 1.14 times, in *Leccinum* (Lat.) – by 1.12 times, in *Russula Pers.* (Lat.) – by 1.16 times, in *Boletus edulis* (Lat.) – by 1.09 times, in *Amanita rubescens* (Lat.) – by 1.12 times, in *Leccinum scabrum* (Lat.) – by 1.13 times, in *Leccinum aurantiacum* (Lat.) – by 1.1 times and in *Armillaria mellea* (Lat.) – by 1.11 times.

In Option 2 the concentration of lead in mushrooms decreased from 1.8 to 2.0 times, in particular, in *Cantharellus cibarius* (Lat.) – by 1.9 times, in *Gyroporus cyanescens* (Lat.) – by 2.0 times, in *Laetiporus sulphureus* (Lat.) – by 1.8 times, in *Butyriboletus regius* (Lat.) – by 1.84 times, in *Leccinum* (Lat.) – by 1.86 times, in *Russula Pers.* (Lat.) – by 1.9 times, in *Boletus edulis* (Lat.) – by 1.91 times, in *Amanita rubescens* (Lat.) – by 1.8 times, in *Leccinum scabrum* (Lat.) – by 1.85 times, in *Leccinum aurantiacum* (Lat.) – by 1.83 times and in *Armillaria mellea* (Lat.) – by 1.81 times.

In Option 3 the concentration of lead in mushrooms decreased from 1.22 to 1.28 times, in particular, in *Cantharellus cibarius* (Lat.) – by 1.23 times, in *Gyroporus cyanescens* (Lat.) – by 1.29 times, in *Laetiporus sulphureus* (Lat.) – by 1.22 times, in *Butyriboletus regius* (Lat.) – by 1.26 times, in *Leccinum* (Lat.) – by 1.27 times, in *Russula Pers.* (Lat.) – by 1.23 times, in *Boletus edulis* (Lat.) – by 1.27 times, in *Amanita rubescens* (Lat.) – by 1.28 times, in *Leccinum scabrum* (Lat.) – by 1.23 times, in *Leccinum aurantiacum* (Lat.) – by 1.22 times and in *Armillaria mellea* (Lat.) – by 1.26 times (Fig. 3).




Fig. 3. Concentration of lead in mushrooms

The source is based on our own research findings

In Option 1 (Table 4) the concentration of cadmium in mushrooms decreased from 1.08 to 1.2 times, in particular, in *Cantharellus cibarius* (Lat.) – by 1.2 times, in *Gyroporus cyanescens* (Lat.) – by 1.14 times, in *Laetiporus sulphureus* (Lat.) – 1.15 times, in *Butyriboletus regius* (Lat.) – by 1.16 times, in *Leccinum* (Lat.) – by 1.13

Table 4

Cadmium concentration in mushrooms, mg/kg

Kind of mushrooms	MPC	The control	Option 1	Option 2	Option 3
<i>Cantharellus cibarius</i> (Lat.)	0.1	0.06	0.05	0.03	0.04
<i>Gyroporus cyanescens</i> (Lat.)	0.1	0.16	0.14	0.09	0.12
<i>Laetiporus sulphureus</i> (Lat.)	0.1	0.15	0.13	0.08	0.12
<i>Butyriboletus regius</i> (Lat.)	0.1	0.14	0.12	0.07	0.11
<i>Leccinum</i> (Lat.)	0.1	0.17	0.15	0.09	0.13
<i>Russula Pers.</i> (Lat.)	0.1	0.65	0.59	0.37	0.53
<i>Boletus edulis</i> (Lat.)	0.1	0.17	0.15	0.09	0.14
<i>Amanita rubescens</i> (Lat.)	0.1	0.15	0.13	0.08	0.12
<i>Leccinum scabrum</i> (Lat.)	0.1	0.17	0.15	0.09	0.14
<i>Leccinum aurantiacum</i> (Lat.)	0.1	0.13	0.12	0.07	0.10
<i>Armillaria mellea</i> (Lat.)	0.1	0.17	0.15	0.09	0.13

The source is based on our own research findings

times, in *Russula Pers.* (Lat.) – by 1.1 times, in *Boletus edulis* (Lat.) – by 1.13 times, in *Amanita rubescens* (Lat.) – by 1.15 times, in *Leccinum scabrum* (Lat.) – by 1.13 times, in *Leccinum aurantiacum* (Lat.) – by 1.08 times and in *Armillaria mellea* (Lat.) – by 1.13 times.

In Option 2 the concentration of cadmium in mushrooms decreased from 1.75 to 2.0 times, in particular, in *Cantharellus cibarius* (Lat.) – by 2.0 times, in *Gyroporus cyanescens* (Lat.) – by 1.77 times, in *Laetiporus sulphureus* (Lat.) – by 1.87 times, in *Butyriboletus regius* (Lat.) – by 2.0 times, in *Leccinum* (Lat.) – by 1.88 times, in *Russula Pers.* (Lat.) – by 1.75 times, in *Boletus edulis* (Lat.) – by 1.88 times, in *Amanita rubescens* (Lat.) – by 1.87 times, in *Leccinum scabrum* (Lat.) – by 1.88 times, in *Leccinum aurantiacum* (Lat.) – by 1.85 times and in *Armillaria mellea* (Lat.) – by 1.88 times.

In Option 3 the concentration of cadmium in mushrooms decreased from 1.21 to 1.5 times, in particular, in *Cantharellus cibarius* (Lat.) – by 1.5 times, in *Gyroporus cyanescens* (Lat.) – by 1.33 times, in *Laetiporus sulphureus* (Lat.) – by 1.25 times, in *Butyriboletus regius* (Lat.) – by 1.27 times, in *Leccinum* (Lat.) – by 1.3 times, in *Russula Pers.* (Lat.) – by 1.22 times, in *Boletus edulis* (Lat.) – by 1.21 times, in *Amanita rubescens* (Lat.) – by 1.25 times, in *Leccinum scabrum* (Lat.) – by 1.21 times, in *Leccinum aurantiacum* (Lat.) – by 1.3 times and in *Armillaria mellea* (Lat.) – by 1.3 times (Fig. 4).




Fig. 4. Concentration of cadmium in mushrooms
The source is based on our own research findings

Conclusions and prospects for further research. Summarizing the obtained results, it should be noted that the most effective way to reduce the concentration of lead and cadmium in mushrooms was the option of soaking and boiling mushrooms in the water without mineral residue. The most effective way to reduce the concentration of zinc and copper was Option 3, where distilled water was used. Thus, the concentration of zinc in mushrooms decreased from 1.01 to 1.04 times, the concentration of copper – from 1.0 to 1.2 times, the concentration of lead – from 1.09 to 1.16 times and the concentration of cadmium – from 1.08 to 1.2 times, when soaking and boiling them in tap water for 3 hours. When soaking and boiling mushrooms in the water without mineral residue, the concentration of zinc decreased from 1.01 to 1.04 times, the concentration of copper – from 1.16 to 1.25 times, the concentration of lead – from 1.8 to 2.0 times and the concentration of cadmium – from 1.75 to 2.0 times. When soaking mushrooms and boiling them in distilled water, the concentration of zinc decreased from 1.13 to 1.18 times, the concentration of copper – from 1.2 to 1.29 times, the concentration of lead – from 1.22 to 1.28 times and the concentration of cadmium – from 1.08 to 1.2 times.

Список використаної літератури

1. Рибалова О.В., Коробкіна К.М. Новий підхід до оцінки забруднення ґрунтів важкими металами. *Proceedings of the II International Scientific and Practical Conference «Topical problems of modern science»*. November 18, 2017. Warsaw, Poland. Vol. 5. p. 86-90.
2. Integrated Risk Information System (IRIS): U.S. Environmental Protection Agency (EPA). URL: <http://www.epa.gov/iris>
3. Калетник Н.Н., Кранов В.П., Орлов А.А. О радиологическом контроле в лесном хозяйстве Украины. *Лесное хозяйство: межсуз. сб. науч. тр.* 1995. № 1. С. 38-39.
4. Краснов В.П. Радіоекологія лісів Полісся України. Житомир : Вид-во «Волинь». 1998. 112 с.
5. Ландін В.П. Сучасна радіаційна ситуація в радіоактивно забруднених лісах України. Проблеми екології лісу і лісокористування на Поліссі України. *Житомир: Вид- во «Волинь»*, 2002. Вип. 4 (10). С. 23-26.
6. Орлов О.О., Краснов В.П., Ландін В.П., Курбет Т.В., Заворотнюк Г.А. Вміст ^{137}Cs у продукції лісового господарства Житомирської області за даними 2002-2004 рр. *Проблеми екології лісів і лісокористування на Поліссі України : зб. наук. праць. Житомир : Вид-во «Волинь»*. 2005. Вип. 5 (11). С. 49-61.
7. Колодій Т.В., Рябчук В.П. Хімічний склад плодових тіл окремих видів роду *Russula s.f. gray*. *Науковий вісник НЛТУ України. Лісове та садово-паркове господарство*, 2011. Вип. 21.4. С. 63-67.
8. Національна доповідь про стан техногенної та природної безпеки в Україні у 2017 році. URL: <http://www.dsns.gov.ua/.../Nacionalna-dopovid-prostan-tehnogennoyi-ta-prirodnoyi-bez>.

9. Український науково-дослідний інститут цивільного захисту України. Статистика. Аналіз масиву карток обліку пожеж. URL: <https://undicz.dsns.gov.ua/ua/Analiz-masivu-kartok-oblikupozhezh.html>.
10. Грабовський О. В., Рошко В.Г., Ніколайчук О.І. Акумуляція важких металів ґрунтом та рослинними об'єктами в умовах антропогенного навантаження. *Наук. вісник УжДУ. Серія Біологія.* Ужгород, 2000. № 8. С. 158–160.
11. Кошкін Е. И. Физиология устойчивости сельскохозяйственных культур: монография. М. : Дрофа, 2010. 638 с.
12. Мельник Н. М., Морозова Т.В. Стан пилку деревних рослин у промислових зонах міста Чернівці. *Наукові основи збереження біотичної різноманітності: тематичний зб. Ін-ту екології Карпат НАН України.* Львів : Ліга-Прес, 2006. Вип. 7. С. 54–60.
13. Регламент максимальних рівнів окремих забруднюючих речовин у харчових продуктах. Державні гігієнічні правила і норми. № 368. ДР-2013 [Чинний від 2013-05-13]. Київ. 2013. 10 с.
14. Врадій О.І. Вплив терміну вимочування у водно-сольовому розчині грибів на концентрацію в них цинку та міді. *Вісник Уманського національного університету садівництва.* 2019. №2. С.75-78.

Список використаної літератури у транслітерації /References

1. Rybalova O.V., Korobkina K.M. (2017). Novyi pidkhid do otsinky zabrudnennia gruntiv vazhkymy metalamy [A new approach to soil contamination by heavy metals]. *Proceedings of the II International Scientific and Practical Conference «Topical problems of modern science».* November 18. Warsaw, Poland. Vol. 5. 86-90. [in Polish].
2. Integrated Risk Information System (IRIS): U.S. Environmental Protection Agency (EPA) (2019). URL: <http://www.epa.gov/iris>. [in USA].
3. Kalenky N.N., Kranov V.P., Orlov A.A. (1995). O radyolohicheskem kontrole v lesnom khoziaistve Ukrayny [On radiological control in the forestry of Ukraine]. *Lesnoe khoziaistvo: mezhvuz. sb. nauch. tr. – Forestry: interuniversity collection of scientific papers.* № 1. 38-39. [in Ukrainian].
4. Krasnov V.P. (1998). Radioekolohiia lisiv Polissia Ukrayny [Radioecology of forests in Polissia of Ukraine]. Zhytomyr : Vyd- vo «Volyn». [in Ukrainian].
5. Landin V.P. (2002). Suchasna radiatsiina sytuatsiia v radioaktyvnno zabrudnenykh lisakh Ukrayny. Problemy ekolohii lisu i lisokorystuvannia na Polissi Ukrayny [Current radiation situation in radioactively contaminated forests of Ukraine. Problems of forest ecology and forest management in Polissia of Ukraine.]. Zhytomyr : Vyd- vo «Volyn» - Zhytomyr: «Volyn» Publishing House, Issue. 4 (10). 23-26. [in Ukrainian].

6. Orlov O.O., Krasnov V.P., Landin V.P., Kurbet T.V. (2005). Vmist 137Cs u produktsii lisovoho hospodarstva Zhytomyrskoi oblasti za danymi 2002-2004 rr. [137Cs content in forestry products of Zhytomyr region according to 2002-2004]. (2005). / Problemy ekologii lisiv i lisokorystuvannia na Polissi Ukraine : zb. nauk. prats. Zhytomyr : Vyd-vo «Volyn» - Problems of forest ecology and forest management in the Polissya of Ukraine: a collection of scientific papers. Zhytomyr: «Volyn» Publishing House. Issue. 5 (11). 49-61. [in Ukrainian].

7. Kolodiy T.V., Riabchuk V.P. (2011). Khimichnyi sklad plodovykh til okremykh vydiv rodu Russula s.f. gray. [The chemical composition of the fruiting bodies of certain species of the genus Russula s.f. gray.]. Naukovyi visnyk NLTU Ukraine. Lisove ta sadovo-parkove hospodarstvo - Scientific Bulletin of NLTU of Ukraine. Forestry and gardening, Issue. 21.4. 63-67. [in Ukrainian].

8. Natsionalna dopovid pro stan tekhnogennoi ta pryrodnoi bezpeky v Ukraini u 2017 rotsi (2017). [National report on the state of technogenic and natural security in Ukraine in 2017]. URL: <http://www.dsns.gov.ua/.../Nacionalna-dopovid-prostan-tehnogennoyi-ta-prirodnoyi-bez.html>. [in Ukrainian].

9. Ukrainskyi naukovo-doslidnyi instytut tsyvilnoho zakhystu Ukraine. Statystyka (2019). Analiz masyvu kartok obliku pozhezh [Ukrainian Research Institute of Civil Protection of Ukraine. Statistics. Analysis of arrays of fire accounting cards]. URL: <https://undicz.dsns.gov.ua/ua/Analiz-masivu-kartok-oblikupozhezh.html>. [in Ukrainian].

10. Hrabovskyi A. V., Roshko V., Nykolaichuk A.Y. (2000). Akkumuliatsiya tiazhel'nykh metallov pochvoi y rastytelnymy ob'ektamy v usloviakh antropogennoi nahruzky [Accumulation of heavy metals by soil and plant objects under anthropogenic loading]. Nauk. vestnyk Uzhhorodskoho. Seryia Byolohiya - Scientific Bulletin of Uzhgorod State University. Series Biology. Uzhgorod, № 8. 158-160. [in Ukrainian].

11. Koshkyn E. Y. (2010). Fyzyolohiya ustoychivosti selskokhoziaistvennykh kultur [Physiology of crop sustainability]: monohrafia. M.: Drofa. [in Russian].

12. Melnyk N. M., Morozova T.V. (2006). Sostoianye pyltsы drevesnykh rastenyi v promyslennyykh zonakh horoda Chernovtsы [State of pollen of woody plants in industrial zones of Chernivtsi city]. Nauchnye osnovы sokhranenyia byotycheskogo raznoobraziya: tematycheskyi sb. Yn-ta ekolohyy Karpat NAN Ukrayny – Scientific Basics of Biodiversity Conservation: Thematic Collection of the Institute of Ecology of the Carpathians of NAS of Ukraine. Lvov Lyha-Press. Issue. 7. 54-60. [in Ukrainian].

13. Rehlament maksymalnykh rivniv okremykh zabrudniuichykh rechovyn u kharchovykh produktakh (2013). [Regulation of maximum levels of certain contaminants in food]. Derzhavni higiienichni pravyla i normy – State hygiene rules and regulations № 368. DR-2013 [Chynnyi vid 2013-05-13]. [in Ukrainian].

14. Vradiy O.I. (2019). Vplyv terminu vymochuvannia u vodno-solovomu rozchyni hrybiv na kontsentratsii v nykh tsynku ta midi [Influence of soaking time in

aqueous salt solution of mushrooms on the concentration of zinc and copper in them]. Visnyk Umanskoho natsionalnoho universytetu sadivnytstva – Bulletin of the Uman National University of Horticulture. №2. 75-78. [in Ukrainian].

АННОТАЦІЯ

АНАЛІЗ ЕФЕКТИВНОСТІ ЗАСТОСУВАННЯ КУЛІНАРНОЇ ОБРОБКИ ГРИБІВ ДЛЯ ЗМЕНШЕННЯ В НИХ КОНЦЕНТРАЦІЇ ВАЖКИХ МЕТАЛІВ

Дослідження концентрації важких металів проводились у науковій агрочімічній лабораторії кафедри екології та охорони навколишнього середовища факультету агрономії та лісівництва Вінницького національного аграрного університету. Концентрації Cu, Zn, Pb та Cd досліджуваних грибів визначали методом атомно-абсорбційної спектрометрії після сухої мінералізації. Дослідження впливу кулінарної обробки на концентрацію важких металів у грибах проводили за такою схемою: 1. Контроль: гриби, очищені та промиті водопровідною водою; 2. Варіант 1: гриби замочують у водопровідній воді на 3 години і варять у ній; 3. Варіант 2: гриби замочують у воді без залишків мінеральних речовин на 3 години і варять у ній; 4. Варіант 3: гриби замочують у дистильованій воді на 3 години і варять у ній. Підсумовуючи отримані результати, слід зазначити, що найефективнішим способом зниження концентрації свинцю та кадмію в грибах був варіант замочування та відварювання грибів у воді без залишків мінеральних речовин. Найефективнішим способом зниження концентрації цинку та міді був варіант 3, де використовувалася дистильована вода. Досліджено вплив замочування та кипіння грибів у різних типах води. При замоченні та кип'ятінні грибів у водопровідній воді протягом 3 годин виявлено, що концентрація цинку в них знижується з 1,01 до 1,04 рази, концентрація міді - з 1,0 до 1,2 рази, концентрація свинцю - з 1,09 до 1,16 рази і концентрація кадмію - від 1,08 до 1,2 рази. При замочуванні та кип'ятінні грибів у воді без залишку мінеральних речовин концентрація цинку зменшилася з 1,01 до 1,04 рази, концентрація міді - з 1,16 до 1,25 рази, концентрація свинцю - з 1,8 до 2,0 рази і концентрація кадмію - від 1,75 до 2,0 рази. При замочуванні грибів і кип'ятінні їх у дистильованій воді концентрація цинку зменшилася з 1,13 до 1,18 разу, концентрація міді - з 1,2 до 1,29 разу, концентрація свинцю - з 1,22 до 1,28 разу та концентрація кадмію - з 1,08 в 1,2 рази.

Ключові слова: гриби, концентрація, цинк, мідь, свинець, кадмій, дистильована вода, вода без мінерального залишку, водопровідна вода, переварювання грибів.

Табл. 4. Рис. 4. Літ. 14.

АННОТАЦИЯ

АНАЛИЗ ЭФФЕКТИВНОСТИ ПРИМЕНЕНИЯ КУЛИНАРНОЙ ОБРАБОТКИ ГРИБОВ ДЛЯ СНИЖЕНИЯ В НИХ КОНЦЕНТРАЦИИ ТЯЖЕЛЫХ МЕТАЛЛОВ

Исследовано влияние вымачивания и переваривания грибов в разных видах воды. Установлено, что в грибах концентрация цинка, меди, свинца и кадмия снизилась от 1,01 до 1,04 раза; от 1,0 до 1,2 раза; от 1,09 до 1,16 раза; от 1,08 до 1,2 раза соответственно при вымачивании и переваривании их в водопроводной воде в течение 3-х часов. При вымачивании и переваривании грибов в воде без минерального остатка концентрация цинка, меди, свинца и кадмия снизилась от 1,01 до 1,04 раза; от 1,16 до 1,25 раза; от 1,8 до 2,0 раза; от 1,75 до 2,0 раза соответственно. При вымачивании грибов и переваривании их в дистиллированной воде концентрация цинка, меди, свинца и кадмия снизилась от 1,13 до 1,18 раза; от 1,2 до 1,29 раза; от 1,22 до 1,28 раза; от 1,08 до 1,2 раза.

Ключевые слова: грибы, концентрация, цинк, медь, свинец, кадмий, дистиллированная вода, вода без минерального остатка, водопроводная вода, переваривания грибов.

Табл. 4. Рис. 4. Літ. 14.

Інформація про автора

Врадій Оксана Ігорівна – асистент кафедри екології та охорони навколишнього середовища Вінницького національного аграрного університету, (21008, м. Вінниця, вул. Сонячна, 3, e-mail: oksanavradii@gmail.com).

Врадий Оксана Игоревна – ассистент кафедры экологии и охраны окружающей среды Винницкого национального аграрного университета, (21008, г. Винница, ул. Солнечная, 3, e-mail: oksanavradii@gmail.com).

Vradiy Oksana Ihorivna – assistant of the Department of Ecology and Environmental Protection, Vinnytsia National Agrarian University (21008, Vinnytsia, 3, Soniachna St., e-mail: oksanavradii@gmail.com).